

# CRS REPORT

|                         |                |                  |
|-------------------------|----------------|------------------|
| Hospital-Institution:   | City, Country: | 1st Surgeon Dr:  |
| Patient Identification: | Date of birth: | Date of Surgery: |

## 1. Surgery Data


|                     |                                                                                                                                 |                                                           |                    |                             |
|---------------------|---------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------|--------------------|-----------------------------|
| Procedure title:    | Primary/recurrent:                                                                                                              | Previous treatment:<br><i>Details previous treatment:</i> | Type of procedure: | Markers:                    |
| Type of tumor:      |                                                                                                                                 |                                                           |                    |                             |
| Suspected stage IV? | <i>If yes:</i> Pleura Lung Skin Extraabdominal lymph nodes Liver<br>Abdominal wall Parenchyma Spleen parenchyma<br>Other sites: |                                                           |                    | Performance<br>Status-ECOG: |

## 2. Surgical Approach and Findings


|                                 |                       |                  |                         |              |                        |
|---------------------------------|-----------------------|------------------|-------------------------|--------------|------------------------|
| Approach:                       | <i>If laparotomy:</i> | <i>If ports:</i> | Type of procedures:     | Adhesiolysis | Volume of ascites:     |
| <i>Abdominal wall resection</i> | Scar                  | Ports site       | Rectus abdominis muscle | Right Left   | Posterior sheath Other |

## PERITONEAL CANCER INDEX

PCI of Sugarbaker



**Lesion Size Score**  
LS 0 No tumor seen  
LS 1 Tumor up to 0.5 cm  
LS 2 Tumor up to 5.0 cm  
LS 3 Tumor > 5.0 cm  
or confluence



|                          | PRE   |                   | POST  |                   |
|--------------------------|-------|-------------------|-------|-------------------|
|                          | Score | Not<br>explorable | Score | Not<br>explorable |
| 0 – Central              |       |                   |       |                   |
| 1 – Right upper quadrant |       |                   |       |                   |
| 2 – Epigastrium          |       |                   |       |                   |
| 3 – Left upper quadrant  |       |                   |       |                   |
| 4 – Left flank           |       |                   |       |                   |
| 5 – Left lower quadrant  |       |                   |       |                   |
| 6 – Pelvis               |       |                   |       |                   |
| 7 – Right lower quadrant |       |                   |       |                   |
| 8 – Right flank          |       |                   |       |                   |
| 9 – Upper jejunum        |       |                   |       |                   |
| 10 – Lower jejunum       |       |                   |       |                   |
| 11 – Upper ileum         |       |                   |       |                   |
| 12 – Lower ileum         |       |                   |       |                   |
| PCI SCORE                |       |                   |       |                   |

## COMPLETENESS OF CYTOREDUCTION

CC score

## ALETTI SCORE

*Automatic calculation of the Aletti score based on the data entered on pages 3 and 4*

## HIPEC

Technical:

Protocol:

Duration:

Flow rate:

Temperature:

Particularity/problems:

*If residual disease:*

Diffuse serosal

Mesenteric

Liver

Hepatic hilum

Pancreas

Supradiaphragmatic

Celiac axis

Other location/size of  
residual disease:

# CRS REPORT

## 3. Peritonectomy procedures

|                                                          |                                             |                                                                             |                                                    |                                                          |
|----------------------------------------------------------|---------------------------------------------|-----------------------------------------------------------------------------|----------------------------------------------------|----------------------------------------------------------|
| Anterior-ventral parietal<br>Left Right<br>Partial Total | Capsulectomy<br>Right Left<br>Partial Total | Infracolic greater omentum                                                  | Ascending mesocolic<br>peritonectomy (right)       | Bladder peritonectomy<br>Partial Total                   |
| Paracolic gutter<br>Left Right<br>Partial Total          | Falciform ligament resection                | Supracolic omentectomy with<br>preservation of the gastroepiploic<br>arcade | Transverse mesocolic<br>peritonectomy              | Pouch of Douglas<br>peritonectomy<br>Partial Total       |
| Subphrenic<br>Left Right<br>Partial Total                | Round ligament resection                    | Supracolic omentectomy with<br>resection of the gastroepiploic<br>arcade    | Descending mesocolic<br>peritonectomy (left)       | Pararectal peritonectomy<br>Left Right<br>Partial Total  |
| Foramen of Winslow<br>Ant. wall Post. wall               | Skeletonization of left gastric<br>vessels  | Lesser omentectomy<br>(lesser omentum)                                      | Sigmoid mesocolic<br>peritonectomy                 | Iliac fossa peritonectomy<br>Left Right<br>Partial Total |
| Morrison's pouch<br>Partial Total                        | Hepatoduodenal ligament                     | Subpyloric space<br>peritonectomy                                           | Mesenteric peritonectomy<br>Partial Subtotal Total | Inguinal orifice peritonectomy<br>Left Right             |
| Ligament of Treitz region                                | Gastrosplenic ligament<br>resection         | Total pancreatic peritonectomy<br>Partial Total                             | Other :                                            | Posterior pelvic peritoneum<br>Partial Total             |

## 4. Visceral resections

|                                                                                                                                                                                                                                                                                                                   |                                                                                                                                                                                                                                                                                                                                     |                                                                                                                                                                    |                                                                                                                                                                                                                                                                                                                                                   |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Liver</b><br>Non-anatomical resection<br>(wedge)<br>Segmentectomy:<br>Hepatectomy: Right Left<br>Cholecystectomy<br>Distal pancreatectomy<br>Pancreaticoduodenectomy<br>(Whipple)<br>Splenectomy<br><b>Stomach - gastrectomy</b><br>Wedge resection<br>Distal Subtotal Total<br><b>Duodenum</b><br>D1 D2 D3 D4 | <b>Jejunum</b><br>Proximal Mid. Distal<br><b>Ileum</b><br>Proximal Mid. Distal<br>Appendectomy<br>Cecectomy<br><b>Colectomy</b><br>Right Left<br>Transverse Sigmoid<br>Extended hemicolectomy<br>Right Left<br>Subtotal Total colectomy<br>Anterior resection (high rectum)<br>Low anterior resection<br>Abdominoperineal resection | Nb. of anastomosis:<br>1.<br>Type Mechanism Orientation<br>2.<br>Type Mechanism Orientation<br>Residual small bowel: cm<br>Stoma:<br>Permanent Temporary<br>Other: | Kidney: Right Left Partial<br>Ureters: Right Left Proximal Mid.<br>Distal<br>Bladder resection: Partial Total<br>Urinary reconstruction:<br>Urinary material:<br>Hysterectomy: Total Subtotal<br>Uterine stripping Vaginectomy<br>Salpingo oophorectomy: Right Left<br>Prostatectomy<br>Seminal vesicle: Right Left<br>Spermatic cord: Right Left |
|                                                                                                                                                                                                                                                                                                                   |                                                                                                                                                                                                                                                                                                                                     | Extra-abdominal resection:                                                                                                                                         | Diaphragm tendinous portion: Right Left<br>Muscle resection: Right Left<br>Direct closure Pedicled flap<br>Prosthetic mesh                                                                                                                                                                                                                        |

# CRS REPORT

## 5. Lymphadenectomy

### Pelvic

|                 |       |      |
|-----------------|-------|------|
| External iliac: | Right | Left |
| Internal iliac: | Right | Left |
| Obturator:      | Right | Left |
| Common iliac:   | Right | Left |

### Retroperitoneal (infra-renal)

Paracaval  
Interaortocaval  
Retrocaval  
Para-aortic  
Epigastric vessels  
Retroperitoneal (suprarenal)  
Coeliac axis  
Superior mesenteric artery  
Common bile duct  
Common hepatic artery  
Hepatic hilar  
Greater omentum  
Lesser omentum  
Left gastric artery  
Splenic hilum

|               |       |      |
|---------------|-------|------|
| Para-cardiac: | Right | Left |
|---------------|-------|------|

|          |       |      |
|----------|-------|------|
| Inguinal | Right | Left |
|----------|-------|------|

### Supradiaphragmatic

Subxiphoid  
Cardiophrenic angle: Right Left  
Mediastinal  
Cervical: Right Left  
Axillary: Right Left

## 6. Verification of occult spaces

### Natural occult peritoneal spaces

Superior paraduodenal recess  
Inferior paraduodenal recess  
Right subhepatic space  
Superior recess of the lesser sac  
Subpyloric space  
Hepatocaval space/groove  
Liver fissures

### Difficult peritoneal sites to inspect

Splenic hilum  
Posterior edge of the spleen  
Ileocecal region  
Sigmoid recess  
Small bowel, its mesentery and folds  
Omental attachment along the colon and epiploic appendages  
Right psoas-caval groove and psoas muscle  
Previous surgical planes

Others:

### Peritoneal tunnels

|                                     |                   |
|-------------------------------------|-------------------|
| Inguinal canal (M)                  | Canal of Nuck (F) |
| Tunnel formed by the hepatic bridge |                   |
| Foramen of Winslow                  |                   |

## 7. Other details

Duration of surgery:

| Drain | N°: | Type: | Location: | Comments: |
|-------|-----|-------|-----------|-----------|
|-------|-----|-------|-----------|-----------|

|                    |                       |    |
|--------------------|-----------------------|----|
| N° RBC transfused: | Estimated blood loss: | cc |
|--------------------|-----------------------|----|

**Severe complications** during the operation:

# CRS REPORT

## 8. Remarks